



TITLE:

# A Note on Mountain Pass Solutions for a Class of Allen-Cahn Models (Nonlinear Partial Differential Equations, Dynamical Systems and Their Applications)

AUTHOR(S):

Byeon, Jaeyoung; Rabinowitz, Paul H.

---

CITATION:

Byeon, Jaeyoung ...[et al]. A Note on Mountain Pass Solutions for a Class of Allen-Cahn Models (Nonlinear Partial Differential Equations, Dynamical Systems and Their Applications). 数理解析研究所講究録 2014, 1881: 1-17

ISSUE DATE:

2014-04

URL:

<http://hdl.handle.net/2433/195655>

RIGHT:

# A Note on Mountain Pass Solutions for a Class of Allen-Cahn Models

Jaeyoung Byeon

Department of Mathematical Sciences, KAIST  
291 Daehak-ro, Yuseong-gu, Daejeon 305-701  
Republic of Korea

E-mail : byeon@kaist.ac.kr,

Paul H. Rabinowitz

Department of Mathematics,  
University of Wisconsin-Madison,  
Madison, Wisconsin, 53706, USA  
E-mail: rabinowi@math.wisc.edu

March 16, 2013

## Abstract

For an Allen-Cahn model problem having 0 and 1 as solutions, the authors recently established the existence of a large family of solutions between 0 and 1 [13]. These solutions were obtained as constrained minimizers of a functional associated with the model or as limits of such minimizers. In this note, it is shown that corresponding to each minimizer,  $U$ , there is a second solution,  $V$ , of the problem which is of mountain pass type with  $0 < V < U < 1$ .

## 1 Introduction

In a recent paper [13], the authors studied the following Allen-Cahn model equation:

$$(1.1) \quad -\Delta u + A_\varepsilon(x)G'(u) = 0, \quad x \in \mathbb{R}^n$$

where  $G(u) = u^2(1 - u)^2$  is a double well potential,  $\varepsilon > 0$ , and  $A_\varepsilon(x) = 1 + A(x)/\varepsilon$  with  $0 \leq A \in C^1(\mathbb{R}^n)$ , 1-periodic in  $x_1, \dots, x_n$ ,  $\Omega$  is the support of  $A|_{[0,1]^n}$  and has a smooth boundary, and  $\bar{\Omega} \subset (0, 1)^n$ . Then  $u \equiv 0$  and  $u \equiv 1$  are solutions of (1.1), the so-called pure states, and the question of interest is whether there are other solutions of (1.1) having  $0 < u < 1$ , i.e. so called mixed states. There have been many papers written on this topic, both for Allen-Cahn models: [1]–[4], [15], [18], [20]–[21], as well as in a more general context motivated by Aubry-Mather Theory and initiated by Moser: [17], [8]–[12], [22], [24]. In particular, it was shown in [13] that there is an infinitude of such mixed states that shadow 0 and 1 in any prescribed way on a spatially periodic array of sets. To make this statement more precise, let  $T \subset \mathbb{Z}^n$ , and set

$$A^T = \cup_{i \in T} (i + \Omega); \quad B^T = \cup_{i \in \{\mathbb{Z}^n \setminus T\}} (i + \Omega).$$

Then one of the main results in [13] contains

**Theorem 1.2.** *For  $G$  and  $A$  as above, there is an  $\varepsilon_0 > 0$  such that for each  $\varepsilon \in (0, \varepsilon_0)$  and each  $T \subset \mathbb{Z}^n$ , there is a solution  $U_{\varepsilon, T}$  of (1.1) with  $0 < U_{\varepsilon, T} < 1$ . Moreover as  $\varepsilon \rightarrow 0$ ,  $U_{\varepsilon, T} \rightarrow 1$  uniformly on  $A^T$  and  $U_{\varepsilon, T} \rightarrow 0$  uniformly on  $B^T$ .*

A more detailed statement can be found in [13]. For finite  $T$ ,  $U_{\varepsilon, T}$  is obtained by a minimization argument where pointwise constraints are imposed on the class of admissible functions. Then, with the aid of further a priori bounds, the case of arbitrary  $T \subset \mathbb{Z}^n$  is obtained from the finite case by passing to a limit.

The solutions for finite  $T$  are local minima of a functional corresponding to (1.1) while 0 and 1 are global minima. Possessing a local minimum such as  $U_{\varepsilon, T}$  which is not a global minimum is essentially the geometric hypothesis of the Mountain Pass Theorem. For the current setting, it suggests the possibility of finding mountain pass solutions for (1.1). The main goal of this note is to show that corresponding to each local minimizing solution,  $U_{\varepsilon, T}$  of (1.1), there is a second solution,  $V_{\varepsilon, T}$ , of mountain pass type with  $0 < V_{\varepsilon, T} < U_{\varepsilon, T}$ . In §2, we will show how to prove this for the case of  $T$  finite. Then in §3, we present a slightly refined result.

There is another way to view the geometric situation just described that may also be fruitful. A well known result of Matano, Theorem 4.4 of [16], states that for equations like

$$(1.3) \quad -\Delta u + f(x, u) = 0$$

on bounded domains in  $\mathbb{R}^n$ , if there is an ordered pair of solutions,  $v < w$  of (1.3) and  $v$  and  $w$  are stable solutions in an appropriate sense for the corresponding parabolic equation

$$(1.4) \quad u_t - \Delta u + f(x, u) = 0,$$

then there exists a solution,  $u$  of (1.3) with  $v < u < w$ . See also Amann [5]-[7] and Sattinger [23]. We will comment further on this point of view in §2.

## 2 The proof for finite $T$

The goal of this section is to show that if  $T$  is finite and  $\varepsilon \in (0, \varepsilon_0)$  is sufficiently small, there is a mountain pass solution,  $V_{\varepsilon, T}$  of (PDE) such that  $0 < V_{\varepsilon, T} < U_{\varepsilon, T}$ . To do this, first the minimization characterization of  $U_{\varepsilon, T}$  in [13] will be recalled. Let  $\mathcal{W}$  the closure of  $C_0^\infty(\mathbb{R}^n)$  under

$$\|u\| \equiv \left( \int_{\mathbb{R}^n} |\nabla u|^2 dx + \int_{[-1,1]^n} u^2 dx \right)^{\frac{1}{2}}.$$

For  $u \in \mathcal{W}$ , let  $L_\varepsilon(u) = \frac{1}{2}|\nabla u|^2 + A_\varepsilon G(u)$  and  $J_\varepsilon(u) = \int_{\mathbb{R}^n} L_\varepsilon(u) dx$ .

Let  $d^* = \frac{1}{2}|\partial\Omega - \partial[0,1]^n|$  so  $d^* > 0$ . Choose any  $d \in (0, d^*)$  such that if

$$\Omega_d \equiv \{x \in \Omega \mid |x - \partial\Omega| > d\},$$

then  $\partial\Omega_d$  is diffeomorphic to  $\partial\Omega$ . Set  $A_T = \mathbb{Z}^n + \Omega_d$  and  $B_T = (\mathbb{Z}^n \setminus T) + \Omega_d$ . Choose constants  $a, b$  such that  $0 < b < \frac{1}{2} < a < 1$  and set

$$\Gamma(T) = \{u \in \mathcal{W} \mid u \geq a \text{ on } A_T \text{ and } u \leq b \text{ on } B_T\}.$$

Define

$$c_\varepsilon(T) = \inf_{u \in \Gamma(T)} J_\varepsilon(u).$$

Let  $\chi_S$  denote the characteristic function of the set  $S$ . Then from [13], we have:

**Theorem 2.1.** *Let  $A_\varepsilon$  and  $G$  be as above. Then there exists an  $\varepsilon_0 > 0$  and independent of  $T$  with the property that for each  $\varepsilon \in (0, \varepsilon_0)$  and each finite  $T \subset \mathbb{Z}^n$ ,*

$$1^\circ \mathcal{M}_\varepsilon(T) \equiv \{u \in \Gamma(T) \mid J_\varepsilon(u) = c_\varepsilon(T)\} \neq \emptyset.$$

- 2° Any  $U \in \mathcal{M}_\varepsilon(T)$  satisfies  $0 < U < 1$  and is a classical solution of (PDE).
- 3°  $\mathcal{M}_\varepsilon(T)$  is an ordered set:  $U, V \in \mathcal{M}_\varepsilon(T)$  implies  $U < V$ ,  $U > V$ , or  $U \equiv V$ .
- 4° If  $T \subset S \subset \mathbb{Z}^n$ ,  $U_{\varepsilon,T} \leq U_{\varepsilon,S}$  with strict inequality if  $T \neq S$ .
- 5° There exist constants  $C, c > 0$ , independent of  $T$  and of  $\varepsilon \in (0, \varepsilon_0)$ , satisfying

$$|U_{\varepsilon,T}(x) - \chi_{T+[0,1]^n}(x)| \leq C \exp(-cd(x, T)), \quad x \in \mathbb{R}^n$$

where  $d(x, T)$  denotes the distance from  $x$  to  $T$ .

Let  $T \subset \mathbb{Z}^n$  be finite,  $\varepsilon \in (0, \varepsilon_0)$ , and  $U_{\varepsilon,T} \in \mathcal{M}_\varepsilon(T)$ . Then, it is straightforward to show that

$$(2.2) \quad \limsup_{\varepsilon \rightarrow 0} c_\varepsilon(T) = \limsup_{\varepsilon \rightarrow 0} J_\varepsilon(U_{\varepsilon,T}) < \infty.$$

To obtain a mountain pass solution of (1.1) corresponding to  $T$ , set

$$\mathcal{G}_{\varepsilon,T} \equiv \{g \in C([0, 1], W^{1,2}(\mathbb{R}^n)) \mid 0 \leq g(\theta) \leq U_{\varepsilon,T}, g(0) = 0, g(1) = U_{\varepsilon,T}\}$$

and define

$$(2.3) \quad b_\varepsilon(T) = \inf_{g \in \mathcal{G}_{\varepsilon,T}} \max_{\theta \in [0,1]} J_\varepsilon(g(\theta)).$$

We will show:

**Proposition 2.4.**  $\liminf_{\varepsilon \rightarrow 0} \sqrt{\varepsilon} b_\varepsilon(T) > 0$ .

**Proof:** Arguing indirectly, if the result is false, there exists a sequence  $\{\varepsilon_m\}_{m=1}^\infty \subset (0, 1)$  such that

$$\lim_{m \rightarrow \infty} \varepsilon_m = \lim_{m \rightarrow \infty} \sqrt{\varepsilon_m} b_{\varepsilon_m}(T) = 0.$$

For notational convenience, we assume that

$$(2.5) \quad \lim_{\varepsilon \rightarrow 0} \sqrt{\varepsilon} b_\varepsilon(T) = 0.$$

The definition of  $b_\varepsilon(T)$  implies there is a  $g \in \mathcal{G}_{\varepsilon,T}$  such that

$$(2.6) \quad b_\varepsilon(T) \leq \max_{\theta \in [0,1]} J_\varepsilon(g(\theta)) \leq b_\varepsilon(T) + 1.$$

By (2.6) for each  $\theta \in [0, 1]$ ,

$$(2.7) \quad \int_{\mathbb{R}^n} A_\varepsilon G(g(\theta)) \, dx \leq b_\varepsilon(T) + 1.$$

Let  $B_r(x)$  denote an open ball of radius  $r$  about  $x \in \mathbb{R}^n$ . Note that  $b_\varepsilon(T) = b_\varepsilon(T + p)$  for any  $p \in \mathbb{Z}^n$ . By shifting  $T$  by such a  $p$  if need be, we can assume  $0 \in T$ . Suppose  $B_r(z) \subset \Omega_d$ . Since there is a constant,  $\omega_d > 0$  such that for  $x \in \Omega_d$ ,  $A(x) \geq \omega_d$ , it follows that  $A_\varepsilon(x) \geq \omega_d/\varepsilon$ , and by (2.7),

$$(2.8) \quad \begin{aligned} \int_{B_r(z)} G(g(\theta)) \, dx &\leq \int_{\Omega_d} G(g(\theta)) \, dx \leq \int_{\Omega_d} \frac{\varepsilon}{\omega_d} A_\varepsilon(x) G(g(\theta)) \, dx \\ &\leq \frac{\varepsilon}{\omega_d} (b_\varepsilon(T) + 1) \equiv M(\varepsilon) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Set

$$\mathcal{A}_\varepsilon(\theta) = \{x \in B_r(z) \mid g(\theta) > a\}, \quad \mathcal{B}_\varepsilon(\theta) = \{x \in B_r(z) \mid g(\theta) \leq b\},$$

and

$$\mathcal{D}_\varepsilon(\theta) = \{x \in B_r(z) \mid b < g(\theta) \leq a\}.$$

The measure of a set,  $S$ , will be denoted by  $|S|$ . Let  $\alpha(\theta) = |\mathcal{A}_\varepsilon(\theta)|$ ,  $\beta(\theta) = |\mathcal{B}_\varepsilon(\theta)|$ , and  $\delta(\theta) = |\mathcal{D}_\varepsilon(\theta)|$ . Then

$$(2.9) \quad \alpha(\theta) + \beta(\theta) + \delta(\theta) = |B_r(z)| = |B_r(0)|.$$

There is a constant  $\sigma = \sigma(d, a, b) > 0$  such that  $G(g(\theta)(x)) \geq \sigma$  for  $x \in \mathcal{D}_\varepsilon(\theta) \subset \Omega_d$  for  $\theta \in [0, 1]$ . Hence by (2.8), for all  $\theta \in [0, 1]$ ,

$$\sigma \delta(\theta) \leq \int_{\mathcal{D}_\varepsilon(\theta)} G(g(\theta)) \, dx \leq \int_{B_r(z)} G(g(\theta)) \, dx \leq M(\varepsilon)$$

or

$$(2.10) \quad \delta(\theta) \leq M(\varepsilon)/\sigma.$$

Note that by the choice of  $B_r(z)$ ,

$$(2.11) \quad \begin{cases} \alpha(0) = 0, & \beta(0) = |B_r(0)| \\ \alpha(1) = |B_r(0)|, & \beta(1) = 0. \end{cases}$$

We claim that for all  $\varepsilon \in (0, \varepsilon_0)$ , there is a  $\rho = \rho(\varepsilon) > 0$  such that whenever  $|t - s| \leq \rho$ ,

$$(2.12) \quad |\alpha(t) - \alpha(s)| \leq 2M(\varepsilon)/\sigma.$$

Otherwise for some  $\varepsilon \in (0, \varepsilon_0)$  and for each  $\rho$ , there is a  $t = t(\rho), s = s(\rho)$  with  $|t - s| \leq \rho$  and  $|\alpha(t) - \alpha(s)| > 2M(\varepsilon)/\sigma$ . Without loss of generality, it can be assumed that  $\alpha(t) \geq \alpha(s)$ . Therefore

$$(2.13) \quad \alpha(t) > \alpha(s) + 2M(\varepsilon)/\sigma.$$

Set

$$X = \{x \in B_r(z) \mid |g(t)(x) - g(s)(x)| \geq a - b\}.$$

Then

$$(2.14) \quad |X|(a - b)^2 \leq \int_X |g(t)(x) - g(s)(x)|^2 dx \leq \|g(t) - g(s)\|_{W^{1,2}(\mathbb{R}^n)}^2$$

and the right hand side of (2.14) goes to 0 as  $\rho \rightarrow 0$ . On the other hand, by (2.13),

$$|(\mathcal{B}_\varepsilon(s) \cup \mathcal{D}_\varepsilon(s)) \cap \mathcal{A}_\varepsilon(t)| \geq 2M(\varepsilon)/\sigma.$$

Therefore by (2.10), we see

$$|\mathcal{B}_\varepsilon(s) \cap \mathcal{A}_\varepsilon(t)| \geq M(\varepsilon)/\sigma.$$

and this implies

$$(2.15) \quad |X| \geq M(\varepsilon)/\sigma.$$

For small  $\rho$ , (2.15) is contrary to (2.14). Thus (2.12) holds.

In a similar fashion, by making  $\rho$  still smaller if necessary, it can be assumed that if  $|t - s| \leq \rho(\varepsilon)$ ,

$$(2.16) \quad |\beta(t) - \beta(s)| \leq 2M(\varepsilon)/\sigma.$$

Further requiring that  $\varepsilon$  is small compared to  $|B_r(0)|$ , the inequalities (2.12), (2.16), and (2.11) imply there is a  $t^* = t^*(\varepsilon)$  such that  $\alpha(t^*)$  and  $\beta(t^*)$  belong to  $(|B_r(0)|/3, 2|B_r(0)|/3)$ . By Lemma 3.5 in Chapter 2 of [14], for a constant,  $C$ , independent of  $\varepsilon, r, a, b$ , we have

$$\begin{aligned}
 (2.17) \quad & \left(\frac{1}{3}|B_r(0)|\right)^{1-\frac{1}{n}}(a-b) \leq \alpha(t^*)^{1-\frac{1}{n}}(a-b) \\
 & \leq \frac{Cr^n}{|B_r(0) \setminus \{x \in B_r(0) \mid g(t^*)(x) > b\}|} \int_{\mathcal{D}_\varepsilon(t^*)} |\nabla g(t^*)| \, dx \\
 & \leq \frac{Cr^n |\delta(t^*)|^{\frac{1}{2}}}{\beta(t^*)} \left( \int_{\mathcal{D}_\varepsilon(t^*)} |\nabla g(t^*)|^2 \, dx \right)^{\frac{1}{2}} \\
 & \leq \frac{3Cr^n (M(\varepsilon)/\sigma)^{\frac{1}{2}}}{|B_r(0)|} (2J_\varepsilon(g(t^*)))^{\frac{1}{2}} \leq \frac{3\sqrt{2}C}{\sqrt{\sigma\omega_d}|B_1(0)|} \sqrt{\varepsilon}(b_\varepsilon + 1)
 \end{aligned}$$

via (2.10), (2.8) and (2.6). But as  $\varepsilon \rightarrow 0$ , the right hand side of (2.17) approaches 0 while the left hand side is bounded away from 0, a contradiction. This proves Proposition 2.4.  $\square$

Next we have the an upper estimate for  $b_\varepsilon(T)$ .

**Proposition 2.18.**  $\limsup_{\varepsilon \rightarrow 0} \sqrt{\varepsilon} b_\varepsilon(T) < \infty$ .

**Proof:** First, let  $\psi : [0, 1] \rightarrow \mathbb{R}$  be a smooth function such that  $\lim_{s \rightarrow 0} \psi(s) = -\infty$ ,  $\lim_{s \rightarrow 1} \psi(s) = \infty$  and  $\psi$  is monotone increasing. Second, choose a function  $\phi \in C^1(\mathbb{R})$  such that  $0 \leq \phi(s) \leq 1$ ,  $|\phi'(s)| \leq 2$  for  $s \in \mathbb{R}$ ,  $\phi(s) = 0$  for  $s \leq 0$   $\phi(s) = 1$  for  $s \geq 1$ . Now define

$$g(\theta)(x_1, \dots, x_n) = U_{\varepsilon, T}(x_1, \dots, x_n) \phi\left(\frac{x_1 + \psi(\theta)}{\sqrt{\varepsilon}}\right), \quad \theta \in (0, 1).$$

Then using 5° of Theorem 2.1 shows that  $\lim_{s \rightarrow 0} g(s) = 0$ ,  $\lim_{s \rightarrow 1} g(s) = U_{\varepsilon, T}$ , the limits being taken in  $W^{1,2}(\mathbb{R}^n)$ . Hence defining  $g(0) = 0$  and  $g(1) = U_{\varepsilon, T}$ , shows  $g \in \mathcal{G}_{\varepsilon, T}$  and

$$(2.19) \quad b_\varepsilon(T) \leq \max_{\theta \in [0, 1]} J_\varepsilon(g(\theta)).$$

To estimate the right hand side of (2.19), first observe that

$$\max_{\theta \in [0, 1]} \int_{\mathbb{R}^n} |\nabla g(\theta)|^2 dx \leq 2 \int_{\mathbb{R}^n} (|\nabla U_{\varepsilon, T}|^2 + \frac{1}{\varepsilon} U_{\varepsilon, T}^2 |\varphi'(\frac{x_1 + \psi(\theta)}{\sqrt{\varepsilon}})|^2) dx$$



and

$$\int_{\mathbb{R}^n} U_{\varepsilon,T}^2 |\varphi'(\frac{x_1 + \psi(\theta)}{\sqrt{\varepsilon}})|^2 dx \leq 4 \int_{0 \leq x_1 + \psi(\theta) \leq \sqrt{\varepsilon}} U_{\varepsilon,T}^2 dx$$

Since  $0 \leq U_{\varepsilon,T} \leq 1$ , again by 5° of Theorem 2.1, there exists a constant  $D_1 > 0$ , independent of small  $\varepsilon > 0$  such that for each  $l \in \mathbb{R}$ ,

$$\int_{x_1=l} U_{\varepsilon,T}^2 dx_2 \cdots dx_n \leq D_1.$$

This inequality implies that

$$\int_{0 \leq x_1 + \psi(\theta) \leq \sqrt{\varepsilon}} U_{\varepsilon,T}^2 dx \leq D_1 \sqrt{\varepsilon}.$$

Hence by 5° of Theorem 2.1 yet again, there is a constant,  $D_2$  independent of small  $\varepsilon > 0$ , so that

$$(2.20) \quad \max_{\theta \in [0,1]} \int_{\mathbb{R}^n} |\nabla g(\theta)|^2 dx \leq D_2 / \sqrt{\varepsilon}.$$

Next noting that

$$\begin{aligned} \int_{\mathbb{R}^n} A_{\varepsilon}(x) G(g(\theta)) dx &= \int_{0 \leq x_1 + \psi(\theta) \leq \sqrt{\varepsilon}} A_{\varepsilon}(x) G(g(\theta)) dx \\ &\quad + \int_{\sqrt{\varepsilon} < x_1} A_{\varepsilon}(x) G(U_{\varepsilon,T}) dx \end{aligned}$$

and estimating as above shows there is a constant  $D_3 > 0$ , independent of small  $\varepsilon > 0$  and  $\theta \in [0, 1]$  such that

$$(2.21) \quad \int_{\mathbb{R}^n} A_{\varepsilon}(x) G(g(\theta)) dx \leq D_3 / \sqrt{\varepsilon}.$$

Combining (2.19) - (2.21) then yields the Proposition. □

Now the main result of this section can be stated.

**Theorem 2.22.** *For each finite  $T \subset \mathbb{Z}^n$ , there is an  $\varepsilon_1(T) > 0$  such that for any  $\varepsilon \in (0, \varepsilon_1)$ , there is a solution,  $V_{\varepsilon,T}$  of (1.1) with  $0 < V_{\varepsilon,T} < U_{\varepsilon,T}$  and  $J_{\varepsilon}(V_{\varepsilon,T}) = b_{\varepsilon}(T)$ .*

Before proving Theorem 2.22, some remarks are in order. First to take advantage of some results from [11], it is convenient to replace  $G$  by a new function,  $G^*$  defined by  $G^*|_{[0,1]} = G|_{[0,1]}$  and  $G^*(z)$  is 1-periodic in  $z$ . Thus  $G^* \in C^2(\mathbb{R})$  and is a bounded function. Let

$$L_\varepsilon^*(u) = \frac{1}{2}|\nabla u|^2 + A_\varepsilon G^*(u) \quad \text{and} \quad J_\varepsilon^*(u) = \int_{\mathbb{R}^n} L_\varepsilon^*(u) \, dx.$$

When dealing with functions,  $u$ , where  $0 \leq u \leq 1$ ,  $G^*(u) = G(u)$  so  $J_\varepsilon^*(u) = J_\varepsilon(u)$ . In particular any critical point,  $u$ , of  $J_\varepsilon^*$  with  $0 \leq u \leq 1$  will be a critical point of  $J_\varepsilon$  and a solution of (1.1).

Let

$$\mathcal{S}_T \equiv \{u \in W^{1,2}(\mathbb{R}^n) \mid 0 \leq u \leq U_{\varepsilon,T}\}.$$

Consider the parabolic initial value problem associated with  $G^*$ :

$$(2.23) \quad u_t - \Delta u + A_\varepsilon(x)G_u^*(u) = 0, \quad t > 0, \quad x \in \mathbb{R}^n$$

$$u(x, 0) = \varphi(x) \in \mathcal{S}_T.$$

Let  $\Phi^t(\varphi)(x)$  denote the solution of (2.23). Then we have

**Proposition 2.24.** *1° For each  $\varphi \in \mathcal{S}_T$ , there is a solution,  $\Phi^t(\varphi)$  of (2.23) which exists for all  $t > 0$ .*

*2° For each  $t > 0$ ,  $\Phi^t : \mathcal{S}_T \rightarrow \mathcal{S}_T$ .*

*3° For each  $\varphi \in \mathcal{S}_T$ ,  $J_\varepsilon(\Phi^t(\varphi)) \leq J_\varepsilon(\Phi^s(\varphi))$  if  $t \geq s \geq 0$ .*

*4° For each  $\varphi \in \mathcal{S}_T$ , there is a solution,  $\Psi(\varphi)$ , of (1.1), with  $\Psi(\varphi) \in \mathcal{S}_T$  and a sequence  $(t_k^*) \subset \mathbb{R}$  such that  $t_k^* \rightarrow \infty$  and  $\Phi^{t_k^*}(\varphi) \rightarrow \Psi(\varphi)$  in  $W^{1,2}(\mathbb{R}^n)$  as  $k \rightarrow \infty$ .*

Proposition 2.24 was proved in [11] for a more general class of (bounded) nonlinearities than  $A_\varepsilon(x)G_u^*$ , but with  $\mathbb{R}^n$  replaced by an  $n$ -torus. However one can use the heat kernel as in [11] (which in turn was based on arguments in [19]) to convert (2.23) to an operator equation in  $W^{1,2}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  and then continue to argue as in [11] to get 1° – 4° of the Proposition.

Next we have:

**Proposition 2.25.** *On  $\mathcal{S}_T$ ,  $J_\varepsilon = J_\varepsilon^*$  satisfies the Palais-Smale condition, i.e. any sequence  $(u_k) \subset \mathcal{S}_T$ , with  $J_\varepsilon(u_k)$  bounded and  $J'_\varepsilon(u_k) \rightarrow 0$ , has a convergent subsequence.*

**Proof:** Suppose  $(\varphi_k) \subset \mathcal{S}_T$  is a Palais-Smale sequence, i.e.  $J_\varepsilon(\varphi_k) \leq M$  and  $J'_\varepsilon(\varphi_k) \rightarrow 0$ . Due to the form of  $J_\varepsilon$  and of  $U_{\varepsilon,T}$  (recalling 5<sup>o</sup> of Theorem 2.1),  $(\varphi_k)$  is bounded in  $W^{1,2}(\mathbb{R}^n)$ . Hence there is a  $\varphi \in \mathcal{S}_T$  such that along a subsequence,  $\varphi_k \rightarrow \varphi$  weakly in  $W^{1,2}(\mathbb{R}^n)$  and in  $L^2_{loc}(\mathbb{R}^n)$  as  $k \rightarrow \infty$ . Since  $0 \leq \varphi \leq U_{\varepsilon,T}$ , the tails of  $\varphi$  are exponentially small so in fact along our subsequence,

$$(2.26) \quad \|\varphi_k - \varphi\|_{L^2(\mathbb{R}^n)} \rightarrow 0, \quad k \rightarrow \infty.$$

By (2.26), for any  $\psi \in W^{1,2}(\mathbb{R}^n)$ , as  $k \rightarrow \infty$ ,

$$(2.27) \quad \left| \int_{\mathbb{R}^n} A_\varepsilon(x)(G_u(x, \varphi_k) - G_u(x, \varphi))\psi \, dx \right| \leq K\|\varphi_k - \varphi\|_{L^2(\mathbb{R}^n)}\|\psi\|_{L^2(\mathbb{R}^n)} \rightarrow 0$$

where  $K = \|A_\varepsilon\|_{L^\infty}\|G_{uu}\|_{L^\infty[0,1]}$ . Hence  $J'_\varepsilon(\varphi) = 0$  and

$$(2.28) \quad J'_\varepsilon(\varphi_k)(\varphi_k - \varphi) - J'_\varepsilon(\varphi)(\varphi_k - \varphi) = \int_{\mathbb{R}^n} |\nabla(\varphi_k - \varphi)|^2 + A_\varepsilon(G_u(x, \varphi_k) - G_u(x, \varphi))(\varphi_k - \varphi) \, dx \rightarrow 0.$$

as  $k \rightarrow \infty$ . Thus by (2.27)–(2.28),

$$\int_{\mathbb{R}^n} |\nabla(\varphi_k - \varphi)|^2 \, dx \rightarrow 0$$

as  $k \rightarrow \infty$ . Consequently  $\varphi_k \rightarrow \varphi$  in  $W^{1,2}(\mathbb{R}^n)$ . □

Now we can give the

**Proof of Theorem 2.22:** First we note from Proposition 2.4 and (2.2) that for small  $\varepsilon > 0$ ,  $b_\varepsilon(T) > J_\varepsilon(U_{\varepsilon,T})$ .

Now we follow an argument from [12]. Let  $\delta > 0$  and choose  $g \in \Gamma(T)$  such that

$$(2.29) \quad \max_{\theta \in [0,1]} J_\varepsilon(g(\theta)) \leq b_\varepsilon(T) + \delta.$$

By 1<sup>o</sup> – 2<sup>o</sup> of Proposition 2.24, for each  $\tau > 0$  and  $\theta \in [0, 1]$ ,  $\Phi^\tau(g(\theta)) \in \mathcal{S}_T$ . Therefore there is a  $\theta_\tau \in [0, 1]$  such that

$$(2.30) \quad J_\varepsilon(\Phi^\tau(g(\theta_\tau))) \geq b_\varepsilon(T).$$

Hence a subsequence of the points,  $\{\theta_\tau \mid \tau > 0\}$ , converges to some  $\theta_\infty \in [0, 1]$ . Denote this subsequence by  $\{\theta_{\tau_p} \equiv \mu_p \mid p \in \mathbb{N}\}$ . We claim for all  $\tau > 0$ ,

$$(2.31) \quad J_\varepsilon(\Phi^\tau(g(\theta_\infty))) \geq b_\varepsilon(T).$$

Otherwise for some  $\tau > 0$ ,

$$(2.32) \quad J_\varepsilon(\Phi^\tau(g(\theta_\infty))) < b_\varepsilon(T).$$

Since  $g$  is continuous, by (2.32), for large  $p$ ,

$$(2.33) \quad J_\varepsilon(\Phi^\tau(g(\mu_p))) < b_\varepsilon(T).$$

Thus by 3° of Proposition 2.24, for large  $p$ ,

$$(2.34) \quad J_\varepsilon(\Phi^{\mu_p}(g(\mu_p))) < b_\varepsilon(T),$$

contrary to (2.30). Consequently (2.31) holds.

Next observe that by 4° of Proposition 2.24, there is a sequence  $(q_k) \subset \mathbb{R}$  and a solution,  $v_\delta$  of (1.1) in  $\mathcal{S}_T$  such that  $\Phi^{q_k}(g(\theta_\infty)) \rightarrow v_\delta$  in  $W^{1,2}(\mathbb{R}^n)$ . Moreover by (2.31) and (2.29),

$$(2.35) \quad b_\varepsilon(T) \leq J_\varepsilon(v_\delta) \leq b_\varepsilon(T) + \delta.$$

Letting  $\delta \rightarrow 0$ , Proposition 2.25 shows there is a solution of (1.1),  $V_{\varepsilon,T} \in \mathcal{S}_T$  such that along a subsequence,  $v_\delta \rightarrow V_{\varepsilon,T}$  in  $W^{1,2}(\mathbb{R}^n)$  and  $J_\varepsilon(V_{\varepsilon,T}) = b_\varepsilon(T) > c_\varepsilon(T)$ . This implies that  $0 < V_{\varepsilon,T} < U_{\varepsilon,T}$  and completes the proof.  $\square$

To conclude this section, recall that from 3° of Theorem 2.1,  $\mathcal{M}_\varepsilon(T)$  is an ordered set. Suppose that  $U, U^* \in \mathcal{M}_\varepsilon(T)$  and there are no members of  $\mathcal{M}_\varepsilon(T)$  lying between  $U$  and  $U^*$ . Then  $U, U^*$  is called a gap pair of solutions of (1.1) in  $\mathcal{M}_\varepsilon(T)$  and we have:

**Theorem 2.36.** *If  $U, U^*$  is a gap pair of solutions of (1.1) in  $\mathcal{M}_\varepsilon(T)$ , there exists a solution,  $V$  of (1.1) of mountain pass type with  $U < V < U^*$ . Moreover  $J_\varepsilon(V) = d_\varepsilon(T)$  where*

$$d_\varepsilon(T) = \inf_{h \in \mathcal{H}_{\varepsilon,T}} \max_{\theta \in [0,1]} J_\varepsilon(h(\theta))$$

and

$$\mathcal{H}_{\varepsilon,T} \equiv \{h \in C([0, 1], W^{1,2}(\mathbb{R}^n)) \mid U \leq h(\theta) \leq U^*, h(0) = U, h(1) = U^*\}.$$

**Proof:** Once it is shown that

$$(2.37) \quad d_\varepsilon(T) > c_\varepsilon(T),$$

the proof is essentially the same as that of Theorem 2.22. Therefore we merely verify (2.37). Modifying an argument from [11], let  $\rho \in (0, \|U - U^*\|_{L^2(\mathbb{R}^n)})$  and set

$$P_\rho \equiv \{u \in W^{1,2}(\mathbb{R}^n) \mid U \leq u \leq U^* \text{ and } \|U - u\|_{L^2(\mathbb{R}^n)} = \rho\}.$$

We claim

$$(2.38) \quad \inf_{u \in P_\rho} J_\varepsilon(u) \equiv f_\rho(T) > c_\varepsilon(T).$$

Indeed, suppose that (2.38) has been established. Fix  $\rho \in (0, \|U - U^*\|_{L^2(\mathbb{R}^n)})$ . Since for any  $h \in \mathcal{H}_{\varepsilon,T}$ , there is a corresponding  $\theta_h \in (0, 1)$  such that  $h(\theta_h) \in P_\rho$ , by (2.38),

$$d_\varepsilon(T) \geq f_\rho(T) > c_\varepsilon(T)$$

and (2.37) is established.

Finally to verify (2.38), if it is false, there is a sequence  $(u_k) \subset P_\rho$  such that  $J_\varepsilon(u_k) \rightarrow c_\varepsilon(T)$  as  $k \rightarrow \infty$ . The sequence  $(u_k)$  is bounded in  $W^{1,2}(\mathbb{R}^n)$ . Therefore there is a  $v \in W^{1,2}(\mathbb{R}^n)$  with  $U \leq v \leq U^*$  such that, along a subsequence,  $u_k \rightarrow v$  weakly in  $W^{1,2}(\mathbb{R}^n)$  and in  $L^2_{loc}(\mathbb{R}^n)$  as  $k \rightarrow \infty$ . Since for each  $R > 0$ ,  $\int_{B_R(0)} L_\varepsilon(\cdot) dx$  is weakly lower semicontinuous,

$$\int_{B_R(0)} L_\varepsilon(v) dx \leq \liminf_{k \rightarrow \infty} \int_{B_R(0)} L_\varepsilon(u_k) dx \leq \liminf_{k \rightarrow \infty} J_\varepsilon(u_k) = c_\varepsilon(T).$$

Since this is true for all  $R > 0$ ,

$$(2.39) \quad J_\varepsilon(v) \leq c_\varepsilon(T).$$

But  $U, U^*$  is a gap pair so (2.39) implies  $v = U$  or  $v = U^*$ . This is impossible since

$$(2.40) \quad \|U - v\|_{L^2(\mathbb{R}^n)} = \rho.$$

Indeed, to verify (2.40), observe that

$$(2.41) \quad \rho^2 = \|U - u_k\|_{L^2(\mathbb{R}^n)}^2 = \int_{B_R(0)} |U - u_k|^2 dx + \int_{\mathbb{R}^n \setminus B_R(0)} |U - u_k|^2 dx$$

As  $k \rightarrow \infty$  along our subsequence, the first term on the right in (2.41) converges to

$$\int_{B_R(0)} |U - v|^2 dx$$

while for each  $k \in \mathbb{N}$ ,

$$\int_{\mathbb{R}^n \setminus B_R(0)} |U - u_k|^2 dx \leq \int_{\mathbb{R}^n \setminus B_R(0)} |U - U^*|^2 dx.$$

Letting  $R \rightarrow \infty$  and noting that, by 5° of Theorem 2.1,  $U$  and  $U^*$  each decay exponentially to 0 as  $|x| \rightarrow \infty$ , (2.40) follows.  $\square$

**Remark 2.42.** Theorem 1.2 provides the existence of a solution,  $U_{\varepsilon,T}$ , of (1.1) for any  $T \in \mathbb{Z}^n$  and any  $\varepsilon \in (0, \varepsilon_0)$  with  $\varepsilon_0$  independent of  $T$ . If  $T$  is finite, Theorem 2.22 gives another solution,  $V_{\varepsilon,T}$ , for a more restricted set of values of  $\varepsilon$ :  $0 < \varepsilon < \varepsilon_1(T)$  where  $\varepsilon_1(T)$  is possibly very small. The size restriction on  $\varepsilon_1(T)$  is required to get  $b_\varepsilon(T) > c_\varepsilon(T)$ . It is natural to ask whether the range of values for the existence of a second solution can be broadened, perhaps even to  $\varepsilon_0$ . One possible approach to this question is to use the relationship between Theorem 2.22 and the Theorem of Matano [16] mentioned in the Introduction. The latter result is for bounded domains while the current setting is  $\mathbb{R}^n$ . Ignoring this fact, the Matano result requires stability behavior relative to (1.4) for the solutions  $u = 0$  and  $u = U_{\varepsilon,T}$  of (1.1). The linearized equation of (1.1) about  $u = 0$  is

$$(2.43) \quad -\Delta v + A_\varepsilon(x)v = 0$$

where  $A_\varepsilon \geq 1$ , showing that 0 will be a stable solution of (1.4). What happens for initial data for (1.4) near  $U_{\varepsilon,T}$  is less clear. However, at the formal level, a calculation shows  $\min(U_{\varepsilon,T}, a)$  (where  $a$  is as in  $\Gamma(T)$ ) is a weak subsolution of (1.1), at least for small  $\varepsilon$ , and that may suffice for the stability near  $U_{\varepsilon,T}$ . Whether these heuristics can be used to obtain an existence theorem remains an open question.

### 3 A generalization of Theorem 2.22 and some remarks

In this final section we will prove a version of Theorem 2.22 that contains more qualitative information about solutions of mountain pass type. Let

$S \subset T \subset \mathbb{Z}^n$  with  $T$  finite and  $S \neq T$ . Define

$$G_\varepsilon(S, T) \equiv \{g \in C([0, 1], W^{1,2}(\mathbb{R}^n)) \mid U_{\varepsilon,S} \leq g(\theta) \leq U_{\varepsilon,T} \\ \text{and } g(0) = U_{\varepsilon,S}, g(1) = U_{\varepsilon,T}\}$$

and set

$$b_\varepsilon(S, T) = \inf_{g \in G_\varepsilon(S, T)} \max_{\theta \in [0, 1]} J_\varepsilon(g(\theta)).$$

Then corresponding to Proposition 2.4, we have

**Proposition 3.1.**  $\liminf_{\varepsilon \rightarrow 0} \sqrt{\varepsilon} b_\varepsilon(S, T) > 0$ .

**Proof:** The proof is quite close to that of the earlier case so we will be brief. Choose  $q \in T \setminus S$  and  $r, z$  so that  $B_r(z) \subset q + \Omega_d$ . Then following the previous proof yields the Proposition.  $\square$

Next corresponding to Proposition 2.18, we also have

**Proposition 3.2.**  $\limsup_{\varepsilon \rightarrow 0} \sqrt{\varepsilon} b_\varepsilon(S, T) < \infty$ .

To continue, replace the set  $\mathcal{S}_T$  of §2 by

$$\Psi(S, T) = \{u \in W^{1,2}(\mathbb{R}^n) \mid U_{\varepsilon,S} \leq u \leq U_{\varepsilon,T}\}.$$

Then there are analogues of Proposition 2.24 and Proposition 2.25 corresponding to this replacement. In particular,  $\Psi(S, T)$  is invariant under  $\Phi^t$ . As earlier, these tools and the proof of Theorem 2.22 yield:

**Theorem 3.3.** *Let  $S$  and  $T$  be as above. Then there is an  $\varepsilon_2 = \varepsilon_2(S, T) > 0$  such that for any  $\varepsilon \in (0, \varepsilon_2)$ , there is a solution,  $W_{\varepsilon,S,T}$  of (1.1) with  $U_{\varepsilon,S} < W_{\varepsilon,S,T} < U_{\varepsilon,T}$  and  $J_\varepsilon(W_{\varepsilon,S,T}) = b_\varepsilon(S, T)$ .*

We conclude with two remarks and some open questions.

**Remark 3.4.** Consider a finite set  $S \subset \mathbb{Z}^n$  and an infinite set  $T \subset \mathbb{Z}^n$  with  $S \subset T$ . For each large integer  $m > 0$ ,  $S \subset T_m \equiv T \cap [-m, m]^n$ . Then by Theorem 2.1, there are corresponding solutions  $U_{\varepsilon,S} \in \Gamma(S)$  and  $U_{\varepsilon,T_m} \in \Gamma(T_m)$ . Theorem 3.3 provides an  $\varepsilon_m = \varepsilon_m(S, T_m)$  such that for any  $\varepsilon \in (0, \varepsilon_m)$ , there is a solution,  $W_{\varepsilon,S,T_m}$  of (1.1) with  $U_{\varepsilon,S} < W_{\varepsilon,S,T_m} < U_{\varepsilon,T_m}$  and  $J_\varepsilon(W_{\varepsilon,S,T_m}) = b_\varepsilon(S, T_m)$ . In [13], we showed that there is an  $\varepsilon_0 > 0$ , independent of  $m > 0$  such that for  $\varepsilon \in (0, \varepsilon_0)$  and large integer  $m > 0$ ,  $U_{\varepsilon,T_m}$  is a solution of (1.1), and that as  $m \rightarrow \infty$ ,  $U_{\varepsilon,T_m}$  converges in  $C_{loc}^2(\mathbb{R}^n)$  to a

solution  $U_{\varepsilon,T}$  of (1.1) with the property that  $\lim_{\varepsilon \rightarrow 0} U_{\varepsilon,T}(x) = 1$  uniformly on  $T + \Omega$  and  $\lim_{\varepsilon \rightarrow 0} U_{\varepsilon,T}(x) = 0$  uniformly on  $(\mathbb{Z}^n \setminus T) + \Omega$ . It is an interesting open question as to whether  $\liminf_{m \rightarrow \infty} \varepsilon_m > 0$ . If this is the case, for small  $\varepsilon > 0$ ,  $W_{\varepsilon,S,T_m}$  converges to a solution  $W_{\varepsilon,S,T}$  of (1.1) with  $U_{\varepsilon,S} < W_{\varepsilon,S,T} \leq U_{\varepsilon,T}$ . Hence, it is then also an interesting open question to see whether  $W_{\varepsilon,S,T} < U_{\varepsilon,T}$  or  $W_{\varepsilon,S,T} = U_{\varepsilon,T}$ .

**Acknowledgments** This research of the first author was supported by the Mid-career Researcher Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (MEST) (No. 2010-0014135).

## References

- [1] FRANCESCA ALESSIO, LOUIS JEANJEAN AND PIERO MONTECCHIARI, Stationary layered solutions in  $\mathbb{R}^2$  for a class of non autonomous Allen-Cahn equations, *Calc. Var. Partial Differential Equations*, **11** (2000), 177–202
- [2] FRANCESCA ALESSIO, LOUIS JEANJEAN AND PIERO MONTECCHIARI, Existence of infinitely many stationary layered solutions in  $\mathbb{R}^2$  for a class of periodic Allen-Cahn equations, *Comm. Partial Differential Equations*, **27** (2002), 1537–1574
- [3] FRANCESCA ALESSIO AND PIERO MONTECCHIARI, Entire solutions in  $\mathbb{R}^2$  for a class of Allen-Cahn equations, *ESAIM Control Optim. Calc. Var.*, **11** (2005), 633–672
- [4] FRANCESCA ALESSIO AND PIERO MONTECCHIARI, Multiplicity of entire solutions for a class of almost periodic Allen-Cahn type equations, *Adv. Nonlinear Stud.*, **5** (2005), 515–549
- [5] HERBERT AMANN, On the existence of positive solutions of nonlinear elliptic boundary value problems, *Indiana Univ. Math. J.*, **21** (1971), 125–146
- [6] HERBERT AMANN, Existence of multiple solutions for nonlinear elliptic boundary value problems, *Indiana Univ. Math. J.*, **21** (1972), 925–935



- [7] HERBERT AMANN, Supersolutions, monotone iterations, and stability, *J. Differential Equations*, **21** (1976), 363–377
- [8] VICTOR BANGERT, On minimal laminations of the torus, *Ann. Inst. Poincare Anal. Non Lineaire*, **6** (1989), 95–138
- [9] UGO BESSI, Many solutions of elliptic problems on  $\mathbb{R}^n$  of irrational slope, *Comm. Partial Differential Equations*, **30** (2005), 1773–1804
- [10] UGO BESSI, Slope-changing solutions of elliptic problems on  $\mathbb{R}^n$ , *Non-linear Anal.*, **68** (2008), 3923–3947
- [11] SERGEY BOLOTIN AND PAUL. H. RABINOWITZ, On the multiplicity of periodic solutions of mountain pass type for a class of semilinear PDE's. *J. Fixed Point Theory Appl.*, **2** (2007), 313–331.
- [12] SERGEY BOLOTIN AND PAUL. H. RABINOWITZ, Hybrid mountain pass homoclinic solutions of a class of semilinear elliptic PDEs, to appear in *Ann. Inst. Poincare Anal. Non Lineaire*
- [13] JAEYOUNG BYEON AND PAUL H. RABINOWITZ, On a phase transition model, to appear in *Calc. Var. Partial Differential Equations*
- [14] O. A. LADYZHENSKAYA AND N. N. URAL'TSEVA, *Linear elliptic and Quasilinear Elliptic Equations*, Academic Press, New York and London, (1968)
- [15] RAFAEL DE LA LLAVE AND ENRICO VALDINOCI, Multiplicity results for interfaces of Ginzburg-Landau Allen-Cahn equations in periodic media, *Adv. Math.*, **215** (2007), 379–426
- [16] HIROSHI MATANO, Asymptotic Behavior and Stability of Solutions of Semilinear Diffusion Equations, *Publ. RIMS, Kyoto Univ.*, **15** (1979), 401–454.
- [17] JÜRGEN MOSER, Minimal solutions of a variational problems on a torus, *Ann. Inst. Poincare Anal. Non Lineaire*, **3** (1986), 229–272
- [18] MATTEO NOVAGA AND ENRICO VALDINOCI, Bump solutions for the mesoscopic Allen-Cahn equation in periodic media. *Calc. Var. Partial Differential Equations*, **40** (2011), no. 1-2, 37–49

- [19] AMNON PAZY, Semigroups of Linear Operators and Partial Differential Equations, Springer Applied Mathematical Sciences series **44**, (1983)
- [20] PAUL H. RABINOWITZ AND ED STREDULINSKY, Mixed states for an Allen-Cahn type equation, *Comm. Pure Appl. Math.*, **56** (2003), 1078–1134
- [21] PAUL H. RABINOWITZ AND ED STREDULINSKY, Mixed states for an Allen-Cahn type equation. II, *Calc. Var. Partial Differential Equations*, **21** (2004), 157–207
- [22] PAUL H. RABINOWITZ AND ED STREDULINSKY, Extensions of Moser-Bangert Theory: Locally Minimal Solutions, *Progress in Nonlinear Differential Equations and Their Applications*, 81, Birkhauser, Boston, (2011)
- [23] Sattinger, D. H., Monotone methods in nonlinear elliptic and parabolic boundary value problems, *Indiana Univ. Math. J.*, **21** (1972), 979–1000.
- [24] ENRICO VALDINOCI, Plane-like minimizers in periodic media: jet flows and Ginzburg-Landau-type functionals, *J. Reine Angew. Math.*, **574** (2004), 147–185